van Bree et al.

Application No.: 09/454,711

Page 3

If the Examiner believes a telephone conference would expedite prosecution of this application, please telephone the undersigned at 650-326-2400.


Respectfully submitted,

Joe Liebeschuetz Reg. No. 37,505

TOWNSEND and TOWNSEND and CREW LLP Two Embarcadero Center, 8th Floor San Francisco, California 94111-3834

Tel: (650) 326-2400 Fax: (650) 326-2422 JOL:adm

PA 3175895 v1



VERSION WITH MARKINGS TO SHOW CHANGES MADE

The paragraph beginning on page 4, line 25, has been amended as follows: Fig. 1: A transgene containing acid α-glucosidase cDNA. The αs1-casein exons are represented by open boxes; α-glucosidase cDNA is represented by a shaded box. The αs1casein intron and flanking sequences (SEQ ID NOS:2 and 3) are represented by a thick line. A thin line represents the IgG acceptor site. The transcription initiation site is marked (1^{\rightarrow}) , the translation initiation site (ATG), the stop codon (TAG) and the polyadenylation site (pA).

The paragraph beginning on page 27, line 16, has been amended as follows: As a further demonstration of the authenticity of α -glucosidase produced in the milk, the N-terminal amino acid sequence of the recombinant α-glucosidase produced in the milk of mice was shown to be the same as that of α -glucosidase precursor from human urine as published by Hoefsloot et al., EMBO J. 7:1697-1704 (1988) which starts with AHPGRP (SEO ID NO:1).

PA 3175895 vl